

Alude

FUNGICIDE

Proven Control Against the Toughest Diseases.

A systemic fungicide, Alude™ provides effective control of various plant diseases including black spot, scab and downy mildew. Alude is a proven performer against Phytophthora and Pythium associated with stem and canker blight. Plus, Alude offers multiple application method options and is tank-mix compatible with many pesticides.

MULTIPLE POINTS OF DEFENSE

- . Multi-site mode of action
- True systemic mobility in the plant, moving both up and down in the plant's conductive tissue
- Enhances plant's natural defense mechanisms
- Decreases plant stress

MIXING AND APPLICATION OPTIONS

- Tank-mix compatible with most pesticides
- Full chemigation label

EPA REG. NO.	71962-1-1001
ACTIVE INGR.	mono- and di-potassium salts of phosphorous acid (53.6%)
FORMULATION	liquid
CHEM. FAMILY	phosphite
FRAC NO.	33
REI	4 hours
PACKAGE SIZE	2 x 2.5 gal, 4 x 1 gal
APPLICATION	spray, drench







CAUTION







KEY USES

Annual and perennial flowers
Bedding plants
Foliage plants
Groundcovers
Deciduous and evergreen trees and shrubs
Vegetable transplants

KEY DISEASES CONTROLLED

(See product label for complete list)

Bacterial blight Downy mildew Phytophthora spp. Pythium spp.

APPLICATION RECOMMENDATIONS

Ornamental plants: Make applications before disease development and in conjunction with good cultural management practices. Use higher rate of application when disease pressure is severe. Do not exceed recommended application rates or apply more frequently than stated on label or plant injury may occur. Do not apply to plants that are heat or moisture stressed. Do not apply to plants that are in a state of dormancy.

Bedding plants: Make applications before disease development and in conjunction with good cultural management practices. Use higher rate of application when disease pressure is severe. Do not exceed recommended application rates or apply more frequently than stated on label or plant injury may occur. Do not apply to plants that are heat or moisture stressed.

Conifers and Christmas trees: Do not exceed indicated application rates or apply more frequently than stated on label in order to avoid plant injury. Do not apply to conifers that are moisture or heat stressed. Do not graze livestock in treated areas of conifer nurseries or plantations. Do not feed forage from treated areas of plantations and or nurseries.

RATES:

Spray Application	2-4 tsp/gal
Spray Application	32-64 fl oz/100 gal
Drench Application	0.375-0.75 tsp/gal
Drench Application	6-12 fl oz/100 gal

See label for complete application rates and recommendations.



