



# **PRE- AND POSTEMERGENCE CONTROL OF KEY BROADLEAF WEEDS**

## **HERBICIDE**

## **Product Advantages**

- Provides both pre- and postemergence control of many broadleaf weeds, including glyphosate-resistant species
- Easily tank-mixed with many pre- and post emergence grass and broadleaf herbicides
- Effective in all tillage systems
- Excellent crop safety

Crop Rotation Interval	
Wheat	4 months
Alfalfa, corn, cotton, sorghum, rice, dry beans, oats, peanuts	9 months
Barley	12 months
Potatoes, sweet corn	18 months
Sugar beets, sunflowers	30 months & bioassay

Refer to the FirstRate® herbicide label for additional crop rotation interval restrictions.

## **Application Timing**



• Preplant: Apply up to two weeks prior to planting or preemergence for additional burndown and early season residual control.

Rate: 0.3 to 0.6 ounce per acre

• Postemergence: Tank-mix with glyphosate to attack and prevent hard-to-control and glyphosate-resistant weeds. FirstRate® may be applied any time prior to the R2 (full flower) growth stage of soybeans

Rate: 0.3 to 0.6 ounce per acre (recommended rate of 0.45 ounce)

## **Control Target Weeds**



Lambsquarters



Marestail



Piaweed



Common Ragweed



Giant Ragweed



Morningglory

## **Weed Spectrum**

Burcucumber<sup>1</sup> Morningglory<sup>2</sup> Cocklebur Mustard, wild3 Lambsquarters<sup>2</sup> Nutsedge, yellow1 Mallow, Venice Pigweed<sup>2</sup> Marestail Ragweed, common Marshelder<sup>3</sup> Ragweed, giant

Smartweed, Pennsylvania Sunflower Thistle, Canada<sup>1</sup> Velvetleaf

- Suppression
- Soil-applied
- 3 Post-applied

Load Chart for FirstRate® herbicide								
			Rate per acre					
Acres per	0.3 oz.		0.45 oz.		0.6 oz.			
load	WSP	Bags	WSP	Bags	WSP	Bags		
40	20	2	30	3	40	4		
80	40	4	60	6	80	8		
160	80	8	120	12	160	16		

Note: Each water soluble packet (WSP) is 0.6 ounce. There are 10 WSP per bag. Each box of FirstRate® herbicide contains four bags or 40 WSF

For more information on FirstRate® herbicide, visit www.dowagro.com/usag.