

PET & LIVESTOCK

PEST CONTROL SPRAY

Flea and Tick control on dogs.
Controls named pests on livestock.

ACTIVE INGREDIENT:	
Permethrin	0.25%
OTHER INGREDIENTS:	99.75%
TOTAL:	100.00%

KEEP OUT OF REACH OF CHILDREN

CAUTION

READ ENTIRE LABEL BEFORE EACH USE
SHAKE WELL BEFORE USING

PRECAUTIONARY STATEMENTS Hazards to Humans & Domestic Animals

CAUTION: Wash thoroughly after handling and before smoking or eating. Avoid contamination of feed and foodstuffs. Remove pets and birds and cover fish aquaria before surface applications. Do not use on humans. Do not allow children or pets to contact treated areas until surfaces are dry. Do not use on dogs under twelve (12) weeks old. Consult a veterinarian before using this product on medicated, debilitated, aged, pregnant or nursing animals. Sensitivities may occur after using ANY pesticide product for pets. If signs of sensitivity occur bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

Environmental Hazards

This product is extremely toxic to fish. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Do not reuse container. Wrap container in several layers of newspaper and discard in trash.

IMPORTANT: READ BEFORE USE

By using this product, user or buyer accepts the following condition, disclaimer of warranties and limitations of liability.

CONDITIONS: the Directions for Use of this product are believed to be adequate and should be followed carefully. However, because of manner of use and other factors beyond Southern Agricultural Insecticide's control, it is impossible for Southern Agricultural Insecticide to eliminate all risks associated with the use of this product. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: THERE ARE NO WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, WHICH EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Southern Agricultural Insecticides is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. Southern Agricultural Insecticides disclaims any liability whatsoever for incidental or consequential damages, including, but not limited to, liability arising out of breach of contract, express or implied warranty (including warranties of merchantability and fitness for a particular purpose), tort, negligence, strict liability or otherwise.

LIMITATIONS OF LIABILITY: THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT SOUTHERN AGRICULTURAL INSECTICIDE'S ELECTION, THE REPLACEMENT OF PRODUCT.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Not for use on plants being grown for sale or other commercial use, or for commercial seed production or for research purposes.

SHAKE WELL BEFORE USING

For emergencies concerning this pesticide product (including health concerns, medical emergencies, or pesticide incidents), call the National Pesticide Telecommunications Network at 1-800-858-7378. Have product container or label with you when calling a poison control center or doctor or going for treatment.

INDOOR APPLICATION

HOUSEHOLD PESTS: For spot application. Do not use as a space spray indoors. Ready to use. Affix applicator to container for application. Do not transfer to another container unless it is a proper pesticide applicator/sprayer.

SURFACE SPRAYING: For 60 day control of Cockroaches, Waterbugs, Palmetto Bugs, Scorpions, Millipedes, Centipedes, Sowbugs, Pillbugs, Silverfish, Firebrats, Spiders, Crickets, Clover mites, Cheese mites, Granary weevils, Rice weevils, Confused Flour Beetles, Rust Red Flour Beetles, Drugstore Beetles, Meal Worms, Grain Mites, and Cadelles: Direct the spray into hiding places, cracks and crevices, and behind shelves and drawers. For silverfish, spray bookcases. For Ants, spray trails, nests and points of entry. Allow all treated areas to dry thoroughly before use.

TO CONTROL FLEAS (ADULTS AND LARVAE) AND TICKS (ADULTS AND LARVAE): Thoroughly spray infested areas, pet beds, resting quarters, nearby cracks and crevices, along and behind baseboard, moldings, window and door frames, and entire areas of the floor and floor covering. Fresh bedding should be placed in animal quarters following treatment. Repeat treatment as needed. Allow to dry thoroughly before allowing pets to contact treated areas.

LIVESTOCK PESTS

Ready to use. Affix applicator to container for application. Do not transfer to another container unless it is a proper pesticide applicator/sprayer.

ON LIVESTOCK: To protect cattle (beef and dairy), goats, sheep, hogs and horses from Horn Flies, Face Flies, House Flies, Mosquitoes and Gnats: Apply a light mist sufficient to wet the surface of the hair. For face flies, spray face and head, but do not spray into eyes.

TO CONTROL STABLE FLIES, HORSE FLIES AND DEER FLIES: apply at a rate of 2 ounces per adult animal, sufficient to wet the hair thoroughly. Repeat treatment daily or at intervals necessary to give continued protection.

TO CONTROL BLOODSUCKING LICE: Apply to the infested areas of the animal, using a stiff brush to get the spray to the base of the hair. Repeat every 3 weeks if required.

TO CONTROL POULTRY LICE: Spray roosts, walls and nests or cages thoroughly. This should be followed by spraying over the birds with a fine mist.

TO CONTROL BEDBUGS AND MITES IN POULTRY HOUSES: Spray crevices of roost poles, cracks in walls, and cracks in nests where the Bedbugs and Mites hide.

PET USE APPLIES TO USE ONLY ON DOGS

ON DOGS; TO CONTROL FLEAS, TICKS AND LICE: Ready to use. Affix applicator to container for application. Do not transfer to another container. Start spraying at the tail, moving the dispenser rapidly and making sure that the animal's entire body is covered, including the legs and underbody. While spraying, fluff the hair so that the spray will penetrate to the skin. Make sure spray wets thoroughly, but do not saturate animal. Do not spray into eyes or face. Avoid contact with genitalia. Repeat application every two weeks if necessary.

DO NOT USE ON CATS OR KITTENS.

OUTDOOR PEST CONTROL

TO HELP CONTROL TICKS IN OUTDOOR ACTIVITY AREAS: Ready to use. Affix applicator to container for application. Do not transfer to another container unless it is a proper pesticide applicator/sprayer. Treat a band of vegetation and leaf litter 6 - 10 feet wide and adjacent to activity areas, especially where dense vegetation occurs.

PESTS ON THE OUTDOOR SURFACE OF BUILDINGS: To temporarily control House Flies, Mosquitoes, Cluster Flies, Gnats and Flying Moths. For outdoor use only as an aid in reducing annoyance from these insects.

Dilute before using. Use clean sprayer. **Carefully measure and mix 1 fluid ounce of Home Pest Control and 12 fluid ounces of water (10 ozs. per gallon).** Using dilute mixture, spray pests on outside surfaces of screens, doors, window frames or wherever these insects may enter the room. Also treat surfaces around light fixtures on porches, in garages and other places where these insects alight or congregate. Repeat as necessary. Flush sprayer with clean water after each use.

ORNAMENTAL FLOWER GARDENS

Ready to use. Affix applicator to container for application. Do not transfer to another container unless it is a proper pesticide applicator/sprayer.

For use on **Roses, African violets, asters, azaleas, begonias, camellias, carnations, chrysanthemums, dahlias, delphiniums, dogwoods, English ivy, eunonymus, fuschia, geraniums, crassula, grape vines, Kentia palms, laurels, marigolds, rhododendrons, rubber plants, snapdragons, stocks, wandering jew, zinnias and other ornamentals**

to kill **Whiteflies, Aphids, Lace bugs, Leafminers, Japanese Beetles, Pavement Ants, Clover Mites, Red Mites, Spider Mites, exposed Thrips, and Armyworms.**

Use sufficient spray to obtain full coverage of all foliage, but do not spray to the point of runoff. When spraying plants, use a sweeping motion spraying no closer than 18 inches from the plant. Avoid heavy wetting of tender foliage, young plants and new growth. Do not spray plants when temperatures exceed 90° F. Contact insects directly whenever possible. Repeat treatment as infestation occur.

For information and questions concerning this product please contact us by email at support@southernag.com.

Southern Agricultural Insecticides, Inc.

Palmetto, FL 34107 Phone, NC 28607
E.P.A. Reg. No. 286-100 No. 829-FL-1