



TREATS A  
1,333 SQ. FT.  
LAWN

**PREVENTS AND  
CONTROLS:**

- Black Spot
- Dollar Spot
- Rust
- Brown Patch
- Powdery Mildew
- Leaf Spot
- ... and more!!

# SYSTEMIC FUNGICIDE RTS

## FUNGICIDE

- Prevents and controls major diseases on roses, flowers, lawns, trees, and shrubs
- Concentrate - Just mix with water and spray
- Systemic Action - Can't be washed off by rain after spray has dried
- Broad spectrum disease control on home lawns and ornamentals

**ACTIVE INGREDIENT:**

Propiconazole (CAS No. 60207-90-1) ..... 1.55%

**OTHER INGREDIENTS:** ..... 98.45%

**TOTAL:** ..... 100.00%

EPA Reg. No. 53883-184    EPA Est. No. 53883-TX-002

**KEEP OUT OF REACH OF CHILDREN  
WARNING**

See Inside Booklet for Additional Precautionary Statements

A **CONTROL SOLUTIONS INC.** PRODUCT

EPA 033011/REV A

**NET CONTENTS: 1 QUART**



## FIRST AID

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

<b>IF IN EYES</b>	<ul style="list-style-type: none"><li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li><li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<b>IF SWALLOWED</b>	<ul style="list-style-type: none"><li>• Call a poison control center or doctor immediately for treatment advice.</li><li>• Have person sip a glass of water if able to swallow.</li><li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li><li>• Do not give anything by mouth to an unconscious person.</li></ul>
<b>IF INHALED</b>	<ul style="list-style-type: none"><li>• Move person to fresh air.</li><li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.</li><li>• Call a poison control center or doctor for further treatment advice.</li></ul>
<b>IF ON SKIN OR CLOTHING</b>	<ul style="list-style-type: none"><li>• Take off contaminated clothing.</li><li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>

**Note to Physician:** If ingested, induce emesis or lavage stomach. Treat symptomatically.

## **PRECAUTIONARY STATEMENTS**

### **Hazards to Humans and Domestic Animals**

Warning - Causes substantial, but temporary eye injury. Do not get in eyes. Wear goggles or safety glasses. Harmful if swallowed, inhaled, or absorbed through skin. Avoid contact with skin or clothing. Avoid breathing vapor or spray mist. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

Do not allow people or pets to contact treated areas until spray is dry.

## **ENVIRONMENTAL HAZARDS**

This pesticide is toxic to fish. Do not apply directly to water. Do not contaminate water when disposing of equipment wash.

## **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This product must not be used in the production of agricultural commodities grown for resale.

**READ ALL INSTRUCTIONS INSIDE BEFORE USE.**

**FOLLOW ALL DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL. POOR DISEASE CONTROL OR INJURY TO DESIRABLE PLANTS MAY RESULT FROM IMPROPER USE OF THIS PRODUCT.**

**DO NOT ALLOW PEOPLE OR PETS INTO TREATED AREAS UNTIL SPRAY HAS DRIED.**

## **GENERAL INFORMATION**

Systemic Fungicide RTS is a water-based systemic fungicide which effectively prevents and controls major diseases on roses, flowers, lawns, trees, and shrubs.

Systemic Fungicide RTS is very effective when used for disease prevention or applied at the first sign of disease. Best results occur when treating early in the season. If disease is already present, Systemic Fungicide RTS will not eliminate it.

Systemic Fungicide RTS, when used as directed, is safe to plants and controls the diseases listed on this label. The use on plants or diseases not listed on the label may result in plant injury and poor disease control.

Do not apply this product to African violets, begonias, Boston ferns, or geraniums.

Contact your local County Extension Service if you need additional information about lawn and ornamental diseases.

## **READY-TO-SPRAY DIRECTIONS FOR USE OUTDOORS ON ROSES, FLOWERS, HOME LAWNS, TREES AND SHRUBS**

### **HOW TO USE THE READY-TO-SPRAY SYSTEM**

#### **Connect**

1. Shake well before using.
2. Connect sprayer to hose.
3. Turn on water.

#### **Spray**

1. To begin spraying, point nozzle in the direction you want to spray.
2. [Bend small plastic tab back and] Turn knob [clockwise] to ON position.
3. Spray evenly to the area to be treated. Refer to the [sight gauge] clear view strip (graduated scale) on the side of the container to determine the amount of product sprayed.

#### **Finish**

1. To stop spraying, [turn knob counter clockwise] to OFF position.
2. Turn off water.
3. Relieve water pressure slowly by [bending plastic tab back and] turning knob to ON position until water slows to a drip. Then turn knob back to OFF position.
4. Disconnect sprayer from hose.

### **APPLICATION RATES: ROSES AND FLOWERS**

- Apply product to the point of run off.
- Before watering, allow spray to fully dry.

Plant	Disease	How Much Product To Apply	Comments
Roses	Powdery Mildew	1 Tablespoon (½ fl. oz.) per gal. of water	For best results, thorough coverage is needed. Begin application at bud break. Treat every 10 days.
	Black Spot, Rust	2 tablespoons (1 fl. oz.) per gal. of water	Treat every 10 days. <b>Note:</b> Under severe conditions, 100% control of black spot may not be achieved.
Calendula	Powdery Mildew	1 Tablespoon (½ fl. oz.) per gal. of water	Apply every 30 days.
Chrysanthemums	Ray Blight	2 Tablespoons (1 fl. oz.) per gal. of water	Apply every 7 days.
Delphinium	Powdery Mildew	1 Tablespoon (½ fl. oz.) per gal. of water	Apply every 14 days.
Snap Dragons	Rust	2 Tablespoons (1 fl. oz.) per gal. of water	Apply every 10 days.
Sweet William	Leaf Spot	1 Tablespoon (½ fl. oz.) per gal. of water	Apply every 14 days.

## LAWNS

- Apply the product to the lawn after mowing or allow the area sprayed to dry completely prior to mowing.
- Prior to application, measure the length and width of area to be treated. Length x width = square feet. (For example, an area 30 ft. long and 20 ft. wide is 600 sq. ft.)

## Established Lawns

Disease	How Much Product to Apply	Comments
Brown Patch	5 fl. oz. in 1 gal. of water per 250 sq. ft. of lawn	Repeat treatment every 10 days, if necessary. First application should take place in May or June before disease is present. For southern lawns, make applications at first sign of disease in spring or fall.
Dollar Spot, Powdery Mildew, Rust, Anthracnose	3 fl. oz. in 1 gal. of water per 250 sq. ft. of lawn	Apply every 14 days.
Red Thread, Dichondra Rust on Dichondra Lawns	5 fl. oz. in 1 gal. of water per 250 sq. ft. of lawn	Repeat every 21 days, if needed, when conditions are favorable for disease to occur.
Stripe Smut	3 fl. oz. in 1 gal. of water per 250 sq. ft. of lawn	Apply in early Spring before grass starts to grow or in Fall after grass is dormant.
Take-All Patch	5 fl. oz. in 1 gal. of water per 250 sq. ft. of lawn	Apply twice in the fall (September-October) and twice in the Spring (April-May) For southern lawns, also make applications twice in the Summer (June-August)

## New Seedings

How Much Product to Apply	Comments
5 fl. oz. in 2 gals. of water per 500 sq ft. of lawn	For faster root development and top growth, apply at the 2 to 3-leaf stage of growth.

## TREES AND SHRUBS

- Apply product to the point of run off.
- Before watering plants, allow spray to fully dry.

Plant	Disease	How Much Product to Apply	Comments
Crab Apple <b>Note:</b> Do not use fruit from treated crab apples for food purposes.	Scab, Rust	½ Tablespoon (¼ fl. oz.) per gal. of water	Apply when buds just begin to break open. Scab- apply every 14 days. Rust- make 3 applications, beginning when buds just begin to break open. Apply at 7-day intervals.
	Powdery Mildew	1 ½ Tablespoons (¾ fl. oz.) per gal. of water	Apply every 14 days
Crape Myrtle	Powdery Mildew	3 Tablespoons (1½ fl. oz.) per gal. of water	Apply every 21 days.
Dogwood	Anthraxnose	1 Tablespoon (½ fl. oz.) per gal. of water	Apply every 14-28 days.
Douglas Fir, Shasta Fir	Needle Rust	3 Tablespoons (1½ fl. oz.) per gal. of water	Douglas Fir-apply every 21 days. Shasta Fir-apply in May-June before disease occurs.
Jeffrey Pine	Tip Blight	3 Tablespoons (1½ fl. oz.) per gal. of water	Make the first application in mid-late winter (Nov.-Jan.) Apply every 2 months up to 3 times
Oak, Red Tip Photinia	Leaf Spot	3 Tablespoons (1½ fl. oz.) per gal. of water	Apply every 14 days.

## NONBEARING FRUIT AND NUT TREE

- Nonbearing fruit and nut trees are those trees that will not produce fruit for at least one year after application of this product.
- Apply product to the point of run off.

Plant	Disease	How Much Product to Apply	Comments
Apple	Scab	½ Tablespoon (¼ fl. oz.) per gal. of water	Begin applications when leaves start growing in the Spring and spray every 14 days.
	Rust	½ Tablespoon (¼ fl. oz.) per gal. of water	Make 3 applications beginning when buds just begin to break open. Apply at 7-day intervals.
	Powdery Mildew	1 Tablespoon (½ fl. oz.) per gal. of water	Begin applications in early Spring and spray every 14 days.
Cherry	Powdery Mildew Leaf Spot	1 Tablespoon (½ fl. oz.) per gal. of water	Begin applications in early Spring and spray every 21 days.
Citrus	Greasy Spot	4 Tablespoons (2 fl. oz.) per gal. of water	Apply June-August every 21 days.
Pecan	Scab, Powdery Mildew, Downy Spot, Zonate Leaf Spot, Fungal Leaf Scorch, Brown Spot	2 Tablespoons (1 fl. oz.) per gal. of water	Begin applications at bud break and make 3 applications at 14-day intervals.
Walnut	Anthraxnose	2 Tablespoons (1 fl. oz.) per gal. of water	Apply every 14 days.



## STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**Pesticide Storage:** Store in cool, dry area away from children and domestic animals.

**Pesticide Disposal:** Pesticide, spray mixture or rinse water that cannot be used according to label instructions must be disposed of at or by an approved waste disposal facility.

**Container Disposal: *For containers equal to or less than 5 gallons:*** Non-refillable container. Do not reuse or refill this container. Clean container promptly. Triple rinse (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container  $\frac{1}{4}$  full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill or by other procedures approved by state or local authorities.

## **WARRANTY STATEMENT**

Control Solutions, Inc. warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on such label only when used in accordance with the Directions For Use. It is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Control Solutions, Inc. To the extent allowed by law, Control Solutions, Inc. shall in no event be liable for consequential, special, or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the Buyer. In addition to the foregoing, no purchaser of this product (other than an end user) shall be entitled to any reimbursement for any loss suffered as a result of any suspension or cancellation of the registration for this product by the U.S. Environmental Protection Agency. Except, as expressly provided herein, Control Solutions, Inc. makes no warranties, guarantees, or representations of any kind, either expressed or implied, or by usage of trade, statutory or otherwise, with regard to the product sold, including, but not limited to merchantability, fitness for a particular purpose, use or eligibility of the product for any particular trade usage. The exclusive remedy of any buyer or user of this product for any and all losses, injuries, or damage resulting from or in any way arising from the use, handling, or application of this product, whether in contract, warranty, tort, negligence, strict liability, or otherwise, shall be damages not exceeding the purchase price paid for this product or, at Control Solutions, Inc. election, the replacement of this product.

Buyer assumes all responsibilities for safety and use not in strict accordance with directions.

## NOTES

Control Solutions, Inc.  
5903 Genoa-Red Bluff  
Pasadena, TX 77507-1041  
EPA Reg. No. 53883-184  
EPA Est. No. 53883-TX-002



# SYSTEMIC FUNGICIDE RTS

- Prevents and controls major diseases on roses, flowers, lawns, trees, and shrubs
- Concentrate - Just mix with water and spray
- Systemic Action - Can't be washed off by rain after spray has dried
- Broad spectrum disease control on home lawns and ornamentals

**ACTIVE INGREDIENT:**

Propiconazole (CAS No. 60207-90-1) . . . **1.55%**

**OTHER INGREDIENTS:** . . . . . **98.45%**

**TOTAL:** . . . . . **100.00%**

**EPA Reg. No. 53883-184    EPA Est. No. 53883-TX-002**

**KEEP OUT OF REACH OF CHILDREN  
WARNING**

See Inside Booklet for Additional Precautionary Statements

**CONTROL SOLUTIONS, INC.**

5903 Genoa-Red Bluff  
Pasadena, Texas 77507-1041

**NET CONTENTS: 1 QUART**

EPA 033011/REV A