



Professional Turf™ Products

Dimension® 0.25 G with DG Pro™

FEATURES & USE TIPS:

- Provides preemergent control of crabgrass and other annual grasses and broadleaf weeds as listed.
- Provides early postemergent crabgrass control in limited conditions.
- Consistently sized particles reduce segregation potential
- Small particle formulation - more per square inch for better herbicide distribution.
- Contains DG Pro™ for greater product dispersibility.

This product is a herbicide that provides control of listed annual grasses and broadleaf weeds in established lawns and ornamental turfs, including golf course fairways, roughs and tee boxes.

New York Use Restrictions: In New York State this product may only be used by professional applicators and at no more than 4.6 pounds per 1000 square feet (equal to 0.5 pound active ingredient per acre); and is prohibited from use in Nassau and Suffolk Counties.

ACTIVE INGREDIENTS:

*Dithiopyr.....	0.25%
OTHER INGREDIENTS.....	99.75%
TOTAL.....	100.00%

†CAS No. 97886-45-8

*This product contains 0.125 pounds of active ingredient, dithiopyr, per 50 pound bag.

Product protected by U.S. Patent No. 41,692,184. Other patents pending.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS KEEP OUT OF REACH OF CHILDREN CAUTION

Causes eye irritation. Harmful if inhaled. Avoid contact with eyes or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Prolonged or frequently repeated skin contact while handling this material may cause allergic reaction in some individuals.

FIRST AID

• **If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contacts, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.

• **If inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for treatment advice.

• **If swallowed:** Call a poison control center or doctor for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

• **If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-757-8951 for emergency medical treatment information.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and highly toxic to other aquatic organisms including oysters and shrimp. Use with care when applying to turf areas adjacent to any body of water. Drift and runoff from treated turf may adversely affect aquatic organisms in adjacent aquatic sites. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

Read the entire label before using this product. Use only according to label instructions.

STORAGE AND DISPOSAL

STORAGE: Store this product only in its original container in a dry, cool, secured storage area. Do not contaminate water, food, or feed by storage and disposal. Keep container closed and away from food, feedstuffs, and domestic water supplies.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticide disposal in accordance with applicable Federal, state or local procedures.

CONTAINER DISPOSAL: Do not reuse bag. Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State or local authorities, by burning. If burned, stay out of smoke.

SPILL: In case of spill, sweep up material and dispose of material according to "pesticide disposal" directions above.

Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in interior landscapes, ornamental gardens or parks, or on golf courses or lawns and grounds.

GENERAL INFORMATION FOR TURF USE

This product is a herbicide that provides control of crabgrass and other annual grasses and broadleaf weeds in established lawns and ornamental turfs, including golf course fairways, roughs, and tee boxes and greens. This product will not harm nearby established ornamentals when used according to label directions.

PRODUCT CODE:

APTDG25DM5

Date: 011604

Supersedes: 110703

PRODUCT CHARACTERISTICS:

Product Type:

Dimension® Herbicide

Carrier:

DG Pro™

Particle Size:

SGN: 150

Net Weight:

50 lbs.

Coverage:

14,700 sq. ft. @
medium rate

EPA Reg. No. 9198-213

EPA Est. 9198-OH-1M, 9198-OH-2B, 9198-AL-001A

Underlined letter is first letter used in run code on bag

Dimension® is a registered trademark of Dow AgroSciences L.L.C.; The Andersons logo is a registered trademark of The Andersons Agriservices, Inc., DG Pro™ and Professional Turf™ are trademarks of The Andersons Agriservices, Inc.

*To order, contact your local
Andersons Lawn Products
distributor.*

*For more information, contact
Andersons at our website:
www.andersonslawnproducts.com
or call 1-800-225-2639.*

USE PRECAUTIONS

TURFGRASS SAFETY

This product may be used on seeded, sodded, or sprigged lawns and ornamental turfs that are well-established. The grass must have developed a good root system and a uniform stand, and have received at least two mowings following its seeding, sodding, or sprigging before it can receive its first application of this product. Use of this product on lawns and ornamental turfs that are not well-established, or on those that have been weakened by weather-, pest-, disease-, chemical-, or mechanical-related stress, may increase the chances for turf injury.

This product should only be applied to lawns and ornamental turfs that are composed of the following turfgrass species that have been determined to be tolerant to applications of this product. When applied as directed under the conditions described, the following established turfgrasses are tolerant to this product:

Cool Season Grasses

Bentgrass, creeping* Fescue, tall
Bluegrass, Kentucky Ryegrass, perennial
Fescue, fine**

Warm Season Grasses

Bahiagrass Centipedegrass
Bermudagrass Kikuyugrass
Buffalograss*** St. Augustinegrass
Carpetgrass Zoysiagrass

* Use of this product on certain varieties of creeping bentgrass, such as 'Cohansey', 'Carmen', 'Seaside', and 'Washington' may result in undesirable turfgrass injury. Not all varieties of creeping bentgrass have been tested.

DO NOT apply this product to Colonial bentgrass (*Agrostis tenuis*) varieties.

**Use of this product on certain varieties of fine fescue may result in undesirable turf injury. The following fine fescue varieties have been found to be sensitive to this product: 'Atlantia', 'Banner', 'Beauty', 'Bilgart', 'CF-2', 'Enjoy', 'HF-93', 'Highlight', 'Ivalo', 'Jamestown', 'Koket', 'Majenta', 'Mary', 'Pennlawn', 'Tamara', 'Tatjana', 'Waldorf', and 'Waldina'. Not all varieties of fine fescue have been tested.

***DO NOT use this product on seedling buffalograss in the spring of the first year of establishment until the turfgrass is fully green and has established new roots.

NOTE: It is not possible to evaluate all turfgrass varieties for tolerance to this product. It is suggested that the user evaluate the response of turfgrass varieties to this product on a limited area, at recommended use rates, prior to initiating large-scale use.

RESEEDING, OVERSEEDING, OR SPRIGGING

Reseeding, overseeding, or sprigging of treated areas with this product should be delayed until 12 weeks from the time of application. Reseeding, overseeding or sprigging before 12 weeks after application may prevent establishment of desirable turfgrasses. When reseeding or overseeding, proper cultural practices such as soil cultivation, irrigation and fertilization should be followed. If overseeding with perennial ryegrass in the fall, overseed 8 weeks or later after a single application. For best results, use mechanical or power seeding equipment (slit seeders) designed to give good seed to soil contact.

OTHER USE PRECAUTIONS

The early postemergence crabgrass control described below is limited. Postemergence applications are only effective on crabgrass if applied before crabgrass has displayed its fifth leaf or first tiller.

For best results, cultural practices that disturb the soil, such as core-, spike-, or hydro-aerification, and verticutting, should be done before application of this product. DO NOT apply this product until the grass has recovered from these cultural practices.

DO NOT use clippings from treated turf for mulching around vegetables or fruit trees.

Keep people and pets off treated areas until dust has settled.

APPLICATION DIRECTIONS

CONTROL OF CRABGRASS

Preemergence and Early Postemergence Control

This product provides "preemergence" control of crabgrass (including the large, smooth, and southern species) when applied prior to the emergence of crabgrass from the ground in established lawns and ornamental turfs.

It can also provide "early postemergence" control of crabgrass. Postemergence crabgrass control will be obtained only when this product is applied prior to the first tillering of crabgrass, which generally corresponds to the time when you can first easily see the crabgrass plants in the lawn or turf. The practical benefit of this product's early postemergence activity is it can give the user a 2-4 week longer period of time to make applications to control crabgrass.

Application Frequency and Timing

This product may be applied as a single application, as a split application, or as a sequential application for crabgrass control in the spring, summer, or fall.

DO NOT apply more than 4.6 lbs. per 1000 sq. ft. per application, and no more than 13.8 lbs. per 1000 sq. ft. per year.

Spring Applications

For single applications made in the spring or early summer, this product should be applied at the appropriate rate listed in Table 1 below. The duration of residual weed control provided by this product is directly related to the total rate applied, but will vary somewhat depending on weather, weed pressure, turfgrass competitiveness, and the user's location within a region. Use the higher recommended rates where weed infestations are dense. Also use the higher rates if applications are made early and longer residual control is desired.

Subsequent, sequential pre- and/or postemergence applications should be made where longer periods of control are desired.

Where split fertilizer applications are recommended, the rates in Table 1 may be split across two applications made 6-10 weeks apart and prior to crabgrass emergence.

Weeds Controlled	Application Rate
Crabgrass: (Large, Smooth, Southern), Foxtail (Green, Yellow)	108 - 160 lbs./acre 2.4 - 3.6 lbs./1,000 sq ft 0.27 - 0.40 lbs. ai/acre
Barnyardgrass	160 - 200 lbs./acre
Chickweed	3.6 - 4.6 lbs./1,000 sq ft
Clover, hop	0.40 - 0.50 lbs. ai/acre
Crabgrass, tropical	
Crowfootgrass	
Dichondra, Carolina	
Lespedeza, common	
Kikuyugrass	
Oxalis	
Buttercup	
Creeping Woodsorrel	
Yellow Woodsorrel	
Pineappleweed	
Ryegrass, annual	
Ryegrass, perennial	
Smutgrass	
Speedwell, com	
Spurge	
Prostrate	
Spotted	

Table 1: (Cont'd.) Recommended Application Rates of This Product

Weeds Controlled	Application Rate
Bluegrass, annual	200 lbs./acre
Bittercress	4.6 lbs./1,000 sq ft
Geranium Carolina	0.50 lbs. ai/acre
Chickweed	
Goosegrass	
Henbit	
Parsley-piert	
Pineappleweed	
Purslane, common	
Sherpardspurse	

Use the higher recommended rates where weed infestations are dense. Also use the higher rates if applications are made early and longer residual control is desired.

Fall Applications

This product can also be applied in the late summer or early fall (late August through November) at the use rates listed in Table 2 to provide control of crabgrass through the early part of the next spring. The fall application should be followed by an appropriate spring application to provide season-long control.

Table 2: Recommended Application Use Rates for Fall Applications (Lbs. of Product/1,000 Square Feet)

Region	Preemergence
North	3.4
Transition	4.6*
South	4.6*
Coastal South	4.6*

*DO NOT apply more than 4.6 lbs. per application, and no more than 13.8 lbs. per year. Postemergence control is limited; see "Crabgrass Control" and "Precautions" sections above.

Tips For Improved Control

For best results, apply this product within a few days after mowing and delay mowing again for a few days after the application.

When treated lawn or ornamental turf areas are watered or receive significant rainfall within a few days after application of this product, improved weed control may result.

Use of split (half-rate) applications spaced 6-8 weeks apart may provide improved weed control.

APPLICATION EQUIPMENT AND INSTRUCTIONS

Apply this product with drop or rotary-type spreaders designed to apply granular herbicides. Avoid the use of spreaders that would apply this product in narrow rows or concentrated bands. Before each application, calibrate the spreader according to the equipment manufacturer's directions for adjusting the spreader settings such that the spreader delivers the appropriate application rate recommended above. Apply this product uniformly over the lawn or ornamental turf area. A more uniform application can be made by spreading half of the required amount over the area and then applying the remaining half at a right angle to the previous direction. Avoid streaking, skips, or overlaps during application. Check equipment frequently to ensure equipment is functioning properly and applying uniform distribution of granules.

GENERAL INFORMATION - LANDSCAPE ORNAMENTALS

This product provides preemergence control of listed annual grass and broadleaf weeds in landscape ornamentals. It may be applied to soil surfaces where ornamental plants are grown for aesthetic purposes in ornamental gardens, and residential properties.

Ornamental plants on this label have shown tolerance to this product when applied as directed. It is impossible, however, to evaluate tolerance for all ornamental plant species or varieties or under all possible growing conditions. Under your growing conditions,

a limited area involving only a few plants should be treated for tolerance evaluation prior to large-scale applications.

USE RATES

Apply this product with a properly calibrated spreader which will assure accurate, even particle distribution. This product may be applied as a single application or as a split application. Refer to the following table for product recommendations when treating a small area.

DO NOT apply more than 4.6 pounds per 1,000 square feet per application and no more than 13.8 pounds per 1,000 square feet per year.

In New York State, this product may only be used by Professional Applicators at no more than 4.6 pounds per 1,000 sq. ft. (0.5 lbs. active ingredient per acre) per year; and is prohibited from use in Nassau and Suffolk Counties.

Lbs of Product per Acre	Lbs of Product per 1,000 Sq Ft	Oz of Product per 100 Sq Ft
200	4.6	7.36

APPLICATION TIMING AND RECOMMENDATIONS

This product is a preemergence herbicide that will not control established weeds. Make applications prior to weed seed germination or to bare ground. The best weed control is obtained when applied to soil that is free of clods, weeds and debris such as leaves. Prior to making an application, control existing vegetation by hand weeding, cultivation or with a postemergence herbicide. Once an application of this product has been made, do not disturb the soil surface as the herbicide barrier will be broken.

PRECAUTIONS

- Only apply this product to established ornamentals.
- Do not apply this product directly to bare root of ornamental plants as injury may result.
- Do not incorporate this product into soil. Dilution of active ingredient and possible injury to plant roots may occur.
- Do not apply to soil around ornamental plants that are under stress due to drought, flooding, excessive fertilizer or soil salts, wind injury, hail, frost damage, winter injury, injury from previously applied pesticides or injury due to insects, nematodes or disease.
- To avoid foliar damage to ornamentals, irrigate immediately after application to remove herbicide particles from the foliage.
- Do not apply under conditions that would affect product distribution. Wind could affect distribution and move product onto sensitive species. Uneven product distribution will result in uneven weed control.
- Do not apply this product on grasses grown for seed.
- Do not graze livestock or feed forage cut from areas treated with this product.

TOLERANT ORNAMENTALS

When applied as directed under the conditions described on this label, ornamentals listed below have shown tolerance in field trials. However, this product has not been tested on all cultivars of each species or under all possible growing condition. Follow directions given above to determine plant tolerance under your growing conditions prior to large scale use.

(continued on page 3)

Common Name	Botanical Name	Tolerant Cultivars
Abelia, Dwarf	Abelia X Grandiflora	Nana
Ajuga	Ajuga reptans	Bronze
	Ajunga genevensis	Bronze Beauty
Almond, Flowering	Prunus glandulosa	
Apple*	Malus pumila	
Arbovitae	Thuja occidentalis	Nigra Pyramidalis Smaragh Techny Woodwardii
Arbovitae, Golden	Thuja orientalis	
Aster, Chinese	Callistephus chinensis	Dwarf Queen
Ash, Green	Fraxinus pennsylvanica	
Ash, Mountain	Sorbus acucuparia	
Ash, Purple	Fraxinus americana	
Azalea	Rhododendron spp.	Brilliant Buccaneer Carror Chimes (Belgian) Exbury Fashion Hardizer Beauty Hershey Red Higasa Hino crimson Holland (Hybrid) Marion Lee Northern Lights Orange Cup Orchid Lights Snow Southern Charm
Azalea, Flame	Rhododendron calendulaceum	
Azalea, Kirishima		
Bamboo, Heavenly		
Barberry	Berberis thunbergii	Aurea Dwarf Pigmy Green Kobold Pygmy Red Rose Glow Atropurpurea Massachusetts
Barberry, Purple		
Bearberry (common)	Arctostaphylos uva-ursi	
Birch, River	Betula nigra	
Blueberry*	Vaccinium spp.	Bluecorp Blue Jay Jersey North Blue Northland
Bottlebrush	Callistemon citrinum	
Boxwood, Japanese		Japonica
Boxwood, Weller	Buxus sempervirens	
Broom	Cytisus spp.	Moonlight
Broom	Genista pilosa	Vancouver Gold
Bugle Carpet		
Carex, Variegated	Carex	
Camellia	Camellia japonica	Debutante Mathotiana Supreme Chansonette
	Camellia sasangua	
Cedar, Red	Juniperus virginiana	
Cockscomb, Plumosa	Celosia cristata	Red Kewpie
Coleus	Coleus blumei	
Copper leaf	Acalypha wilkesiana	
Cotoneaster	Cotonester apiculatus	
Cycads	Cycads revoluta	
Coyotebrush	Baccharis pelularis	
Cypress, Bald	Taxodium distichum	
Cypress, Italian	Cupressus sempervirens	Glauca
Cypress, Leyland	Cupressocyparis leylandii	
Cypress, Japanese False	Chamaecyparis obtusa	Gracilis
Daffodil	Narcissus spp.	King Alfred
Daylily	Hemerocallis spp.	Aztec Gold Bright Yellow (Hybrid) Single Gold (Evergreen) Wilson's Yellow

Common Name	Botanical Name	Tolerant Cultivars
Dogwood	Cornus florida	
Dogwood, American	Cornus sericea	
Douglas Fir	Pseudotsuga menziesii	
Dusty Miller	Senecio cineraria	
Elm, Drake	Ulmus parvifolia	
Euonymus	Euonymus fortunei	Argenteo-variegata Aureo-marginata Colorata Emerald Gaiety Emerald 'N Gold Gold Edge Gold Princess Silver King Tricolor Vegetus
Fan Palm, European	Chamaerops humilis	
Fan Palm, Mexican	Washingtonia robusta	
Fescue	Festuca glauca	
Fetterbush	Leucothoe fontiana	Rainbox
Ficus	Ficus retusa	Nitida
Forsythia	Forsythia X intermedia	Arnold Dwarf Bronxensis Dwarf Lynwood Gold Meadowlark Weeping
Fir Fraser	Abies fraseri	Spring Glory
Gardenia	Gardenia jasminoides	Mystery Radicans
Geranium	Pelargonium X hortorum	
Gum	Eucalyptus citriodora	
Hawthorn	Crataegus spp.	Cockspur White Crimson Cloud Enchantress Jack Evans Washington White Mediterranean Pink Brilliant Hula Girl Blue Bird
Heather, Twisted	Erica cinerea	
Hibiscus	Hibiscus spp.	
Hemlock, Canada	Tsuga canadensis	
Holly	Ilex spp.	Blue Boy Blue Girl
	Ilex X meserveae	Burfordii China Girl Ciomcompacta Forsteri Hellerie Japanese Northern Beauty Needlepoint Nellie R. Stevens Savanah
Holly, Chinese	Ilex X attenuata	
Holly, Japanese	Ilex cornuta	
Holly, Yaupon	Ilex vomitoria	
Honeysuckle	Lonicera japonica	Clavey's Dwarf Halliana Tatarian Canadian White Zebelli Red Hosta Algo Marginata
Hosta	Hosta spp.	
	Hosta lancifolia	
Ice Plant	Carpobrotus edulis	
Impatiens	Impatiens wallerana	
Iris	Iris spp.	Wedgewood Dwarf blue Bulgaria Thorndale
Ivy, English	Hedera helix	
Jasmine, Asian	Trachelopermum asiaticum	
Juniper	Juniperus spp.	Arcadia Armstrong Bar Harbor Blue Chip Blue Pacific Blue Rug Blue Star Broadmoor Buffalo Calgary Carpet

Common Name	Botanical Name	Tolerant Cultivars
		Emerald Sea Emerald Spreader Endora Compacta Fuitlandi Green Gold Tip Hetzl Hughes Manhattan Blue Parsoni Pitzeriana Plumosa Prince of Wales Procumbens Dwarf San Jose Sargent Blude Sargent Green Scandia Scopolorum Moonglow Scopolorum Skyrocket Spartan Tamariscifolia Weberi Youngstown Yukon Belle
Juniperus chinensis		
Juniperus sabina		
King Palm	Archontophoenix cunninghamiana	
Laurel, Australian	Pittosporum tobira	
Laurel, Mountain	Kalmia latifolia	
Leucothoe	Leucothoe fontanesiana	
Lily, African	Agapanthus africanus	Albus Peter Pan
Lily, African Blue		
Lily of the Valley	Pieris japonica	Mt. Fire
Lilyturf	Liriope muscari	Evergreen Giant Lilac Beauty Majestic Monroe White Variegata
Ligustrum, Japanese	Ligustrum japonicum	
Liriope, Green	Liriope spicata	
Maple, Japanese	Acer japonicum	
Maple, Norway	Acer platanoides	
Maple, Red	Acer rubrum	
Maple, Siviler	Acer saccharinum	
Maple, Sugar	Acer saccharum	
Marigold	Tagetes patula	Honeycomb Wheeler's Dwarf Variegata Golden Snowflake Double White
Mock Orange*	Philadelphus spp	
Mondo Grass	Ophiopogon japonicus	
Moss Rose	Portulaca grandiflora	Sunnyside
Myrtle, Crape	Lagerstoemia indica	Faurei Langer Muskogee Standard Pink
Myrtle, Wax	Myrica cerifer	
Nadina	Nadina domestica	Compacta Nana Hardy Red Petite Pink Sister Agnes
Oleander	Nerium oleander	
Oak, Laurel	Quercus laurifolia	
Oak, Pin	Quercus palustris	
Oak, Red	Quercus rubra	
Oak, Southern L	Quercus virginiana	
Oak, Willow	Quercus phellos	
Oteospermum	Osteospermum fruitcosum	Wirligig
Palm, Bangalow	Cortadaria seloana	
Pampas Grass	Bougainvillea glabra	Barbara Karst
Paper Flower	Pachysandra terminalis	
Pachysandra		
Peach*	Prunus persica	
Periwinkle, Dwarf	Vinca minor	
Petunia	Petunia X hybrida	Picoti
Photinia, Red Tip	Photinia X fraseri	
Pieris	Pieris japonica	
Pine, Australian	Pinus nigra	
Pine, Japanese Black	Pinus thunbergiana	
Pine, Loblolly	Pinus teada	
Pine, Longleaf	Pinus palustris	

Common Name	Botanical Name	Tolerant Cultivars
Pine, Mugho	Pinus mugho	
Pine, Scotch	Pinus sylvestris	
Pine, Slash	Pinus eliottii	
Pine, Swiss Mt.	Pinus mugo	
Pine, Virginia	Pinus virginiana	
Pine, White	Pinus strobus	
Pineapple, Guava	Feijoaellowaiiana	
Pittosporum, Japan		
Potentilla	Potentilla fruticosa	Abbotswood
Privet	Ligustrum japonicum	Golden Vicary Regal Texanum Yellow Tipped Wax
Privet, Glossy	Ligustrum lucidum	
Pyracantha	Pyracantha koidzumii	
Queen Pam	Arecastrum rammazzofianum	
Quince, Japanese*		
Rhododendron	Rhododendron spp	Album Cunningham White Fashion Hardy PJM Purple Gem Silvery Pink
Rhododendron, Carolina	Rhododendron carolinianum	
Rhododendron, Catawba	Rhododendron catawbiense	
Rodie Max (Rosebay)	Rhododendron maximum	
Rose	Rosa banksiaae	Luta
Rosemary*	Rosmarinum officinalis	
Rosemary, Bog	Andromeda polifolia	Nana
Salvia	Salvia farinacea	Rhea
Sedum	Sedum spurium	Dragon Blood Red Red Carpet Yellow Fanall
Spiraea	Astilbe X arendsii	Anthony Waterer Red Dolchica Froebel Pink Goldenflame Red Snowmound White Van Houttee White
Spiraea	Spiraea spp.	
Spiraea, Garland	Spiraea spp.	
Spruce, Black Hills	Picea glauca	
Spruce, Colorado Blue	Picea pungens	
Spruce, Norway	Picea abies	
Spruce, White	Picea glauca	
Sourwood	Oxydendrum arboreum	

Common Name	Botanical Name	Tolerant Cultivars
Sweetflag, Grassy-Le		
Sweetgum	Liquidambar styraciflua	
Sycamore	Platanus occidentalis	
Tree Fern (Tiki Fern)	Asparagus virgatus	
Trumpet Flower, Evening	Gelsemium sempervirens	
Tulip	Tulip, spp	Apeldoorn
Verbana, Shrub	Lantana sellowiana	
Verbain	Verbena spp.	St. Paul
Viburnum	Viburnum spp.	American Cranberry Bush Arrowood Common Snowball European Cranberry Bush Linden Mohican Wright
Windmill Palm	Trachycarpus fortunei	
Xylosma	Xylosma congestum	
Yaupon	Ilex vomitoria	Dwarf
Yew	Taxus cuspidata Taxus X media	Denisiformis

*Ornamental species only: DO NOT USE ON FOOD PRODUCING TREES AND ORNAMENTALS

WEEDS CONTROLLED

When used as directed, this product will control or suppress the following annual grass and broadleaf weeds when applied prior to their emergence. This product will not control established grasses, except for crabgrass in early stages of development, and broadleaf weeds; therefore, the area to be treated should be free of weeds prior to application.

GRASSES

Barley
barnyardgrass
bluegrass, annual
brome
crabgrass, large
crabgrass, smooth
crabgrass, Southern
crowfootgrass
dallisgrass (seedling)
goosegrass
foxtail, green
foxtail, yellow
kikuyugrass
oats, wild
ryegrass (annual & perennial)
smutgrass

BROADLEAF WEEDS

bittercress
shepherdspurse
carpetweed
speedwell, corn
chickweed
spurge, garden
geranium, Carolina
spurge, prostrate
henbit
spurge, spotted
knotweed, prostrate
woodsorrel, creeping
lespedeza, common
woodsorrel, yellow
marestail
medic, black
mustard (Brassica spp.)
oxalis, buttercup
pineappleweed
pigweed, redroot
parsley-piert
purslane, common
rockar, London

APPLICATION EQUIPMENT AND INSTRUCTIONS

Apply this product with drop or rotary-type spreaders designed to apply granular herbicides. Avoid the use of spreaders that would apply this product in narrow rows or concentrated bands. Before each application, calibrate the spreader according to the equipment manufacturer's directions for adjusting the spreader settings such that the spreader delivers the appropriate application rate recommended below. Apply this product uniformly over the area to be treated. A more uniform application can be made by spreading half of the required amount over the area and then applying the remaining half at a right angle to the previous direction. Avoid streaking, skips or overlaps during application. Check equipment frequently to ensure equipment is functioning properly and applying uniform distribution of granules.

IMPORTANT: READ BEFORE USE

Read the entire Directions for Use and the Warranty Disclaimer and Limitation of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or Buyer accepts the following Warranty Disclaimer and Limitation of Liability:

WARRANTY DISCLAIMER and LIMITATION of LIABILITY

Manufacturer warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions. Manufacturer makes NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. Where permissible, it is Manufacturer's intent to LIMIT ANY LIABILITY FOR SPECIAL, CONSEQUENTIAL OR INCIDENTAL ECONOMIC DAMAGES to refund of purchase price or replacement of product, at Buyer's choice. Manufacturer DISCLAIMS ANY LIABILITY FOR COMPENSATORY OR OTHER DAMAGES ARISING OUT OF ANY USE CONTRARY TO LABEL DIRECTIONS. Use contrary to label directions is not permitted.

LEGAL RIGHTS

THIS LIMITED WARRANTY GIVES YOU SPECIFIC LEGAL RIGHTS; YOU MAY HAVE OTHER RIGHTS THAT VARY FROM STATE/JURISDICTION TO STATE/JURISDICTION.

APTDG25DM5

Product of USA

Manufactured by:

The Andersons Lawn Fertilizer Division, Inc.

P. O. Box 119, Maumee, OH 43537

1-800-225-2639

Visit us on the web at:

www.andersonsinc.com

Suggested Spreader Settings

These suggested settings are not intended to replace calibration. Please calibrate your spreader before applying product.

50 lbs. treats 20,800 sq. ft. (0.48 acre) at the LOW RATE
50 lbs. treats 14,700 sq. ft. (0.34 acre) at the MEDIUM RATE
50 lbs. treats 10,800 sq. ft. (0.25 acre) at the HIGH RATE

SPREADER SETTINGS

SPREADER	GROUND SPEED	WIDTH OF COVERAGE	LOW RATE	MEDIUM RATE	HIGH RATE
			2.40 lbs./k sq. ft.	3.40 lbs./k sq. ft.	4.60 lbs./k sq. ft.
Andersons Rotaries: AccuPro 1000/2000/SR-2000, (Cone 6)	3 mph	9 feet	I 1/4	J	L 1/4
Andersons SS-2	3 mph	overlap wheels	3 3/4	4 1/4	4 3/4
Scotts Rotaries: R-8, R-8A (Cone 6)	3 mph	9 feet	H 1/2	I 1/2	J 1/2
Earthway Broadcast 2400	2 mph	7 feet	11 1/2	13 1/2	15 1/2
Lely Models WTR, WFR, HR, 1250. (PTO at 450 rpm)	4.5 mph	34 feet	4 1/4 III-C	5 1/4 III-C	6 III-C
LESCO 020093	3 mph	5 feet	D	D 1/2	E
Prizelawn CB85	3 mph	5 feet	3 1/2	3 3/4	4
Vicon (03 Series-Standard Spout)	4.5 mph	30 feet	18	23	29