

AllPro[®] Permethrin ICG

PRECAUTIONARY STATEMENTS Hazards to Humans and Domestic Animals

CAUTION

Harmful if absorbed through skin. Causes moderate eye irritation. Avoid inhalation of dust. Avoid contact with skin, eyes or clothing. Wash thoroughly after handling. Avoid contamination of food and feed.

STOP. READ ENTIRE LABEL BEFORE USE.

FIRST AID

Have the product container or label when calling a poison control center or doctor, or going for treatment.

If on skin or clothing	- Take off contaminated clothing. - Rinse skin immediately with plenty of water for 15-20 minutes. - Call a poison control center or doctor for treatment advice.
If in eyes	- Hold eye open and rinse slowly and gently with water for 15-20 minutes. - Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. - Call a poison control center or doctor for treatment advice.
If inhaled	- Move person to fresh air. - If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. - Call a poison control center or doctor for further treatment advice.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to fish. Do not apply directly to water. Drifts and runoff from treated areas may be hazardous to aquatic organisms in treated areas. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

For outdoor use around the home only.

Three Easy Steps to Use this Product.

Step 1: Determine size of the job, measure the area to be treated.

Step 2: Use spreader setting chart to determine appropriate setting for your spreader.

Step 3: Treat as directed in rate chart. Thorough coverage is important. Water in granules where indicated.

Food utensils such as teaspoons and measuring cups should not be used for food purposes after use with pesticides.

Note: The spreader settings listed below are approximate. Accurately weigh a few pounds into your spreader and apply to a measured area. Then adjust spreader setting if necessary, in order to assure correct rate of application.

SPREADER SETTING NO. (Per 1000 sq. ft.)

Spreader Make & Model	Grass or Dichondra Lawns 2 pounds	St. Augustine Lawn 3 pounds	White grubs, Billbugs 2 1/2 pounds
Central Imperial Make No. 1662-G	6 1/2	7 1/4	7
Central Suburbia Lawn Spreader No 1622-0	6 1/2	7 1/2	7
Cyclone Model B1 Rotary	4 1/2	5	4 3/4
Greenfield Model 105	6 1/2	7 1/4	6 3/4
Ortho Drop Spreader Model 3000	4 1/2	5 1/2	5
Ortho Broadcast Spreader Model 4000	4 1/2	7	6
J. C. Penny Model 6005	7 1/4	8	7 1/2
J.C. Penny Rotary Spreader	5	7	6
Scott Model 35	4	5	4 1/2
Sears Craftsman Model 671.19198	6	7	6 1/2
Spyker 24-22 Rotary	3.6	3.9	3.75
Lesco Rotary	10 (E)	11 (F)	10 (E)
Prizelawn CB Rotary	3.8	4.2	4.0
EarthWay 2400 Rotary	14	15	14
Cyclone Pro	3.25	3.5	3.4
Scotts R8A	F	G	G
Lely Broadcast	2.3	2.5	2.4
Bigfoot	F	G	F-G

Distributed by:
Value Garden Supply
PO Box 585
St. Joseph, MO 64502-0585
(952)884-6477
www.valuegardens.com

EPA Reg. No. 53883-39-769

EPA Est. No. 44616-MO-1@, 44616-GA-1@
Superscript used is first letter of lot number.

Active Ingredient:

Permethrin [3-phenoxyphenyl)methyl (±) cis-trans-3-(2,2-dichloroethyl)-2,2-Dimethyl-
cyclopropanecarboxylate]

By Wt.

0.25%	
99.75%	
100.00%	

*Cis trans isomer ratio: min 35% (±) cis, ,max. 65% (±) trans.

KEEP OUT OF REACH OF CHILDREN CAUTION

DIRECTIONS FOR APPLYING ALLPRO PERMETHRIN ICG

Treatment Areas	Target Pest	Remarks	Use Rate
Band Treatment Around House Foundation	Ants, Fleas, Brown dog ticks, Clover mites, Crickets, Earwigs, Millipedes, Roaches, Sowbugs (Pillbugs), Spiders	These pests are commonly found around or near foundations of houses as well as in lawns. Certain of these pests may enter the houses. Apply a 5-foot band of granules uniformly. Water lightly after application.	3.2 oz. For each 100 sq. ft. area (5 ft. by 20 ft.)
Overall Lawn Application for Grass and Dichondra Lawns	Ants, Armyworms, Brown dog ticks, Chinch bugs, Clover mites, Crickets, Cut worms, Earwigs, Fleas, Lawn moths (sod web-worms), Leafhoppers, Millipedes, Sowbugs Grubs, Japanese beetle grubs, Atranienius sprelulus (dung beetles)	Apply with a fertilizer spreader. See spreader setting chart above. Apply when grass is dry. Water lawn lightly after application. Water grass thoroughly after application. For billbug control, apply when activity is first observed or when chewed or brown grass indicates damage from these insects. For grub control, except dung beetle, treat any time between late July and early October. For dung beetle grub control, apply in spring or early summer when damage is first noticed. Water grass thoroughly with 1/4 - 1/2 inch of water after application. Treat again in late summer or early fall if a second generation occurs.	For application with other types of equipment, apply at the rate of 2 lbs. Per 1000 sq. ft. (an area 20' x 50') or 3.2 oz. Per 100 sq. ft. (an area 10' x10'). For heavy lawns such as St. Augustine grass, the rate should be increased to 3 lbs. Per 1000 sq. ft. (or 4.8 oz. Per 100 sq. ft.) Apply 2.5 lbs. Per 1000 sq. ft. of lawn.
Spot Treatment	Ants Fire Ants	Water lightly after application. Immediately after application, water in thoroughly. A minimum of 1 gallon of water should be applied to each mound area. Apply gently to avoid disturbing ants. Use of high pressure watering equipment may disturb the ants and cause migration, reducing product effectiveness. NOTE: For best results, apply in cool weather, 65° - 80° F, or in early morning or late evening hours. Treat new mounds as they appear.	Apply 1 1/2 teaspoons over and around each ant hill. Apply 1 cup over, and 2 ft. around each fire ant mound.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Storage: Store in cool, dry area away from children and domestic animals.


Disposal: If empty: Do not reuse this container. Do not rinse unless required for recycling. Place in trash.

If partly filled: Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Unless otherwise instructed place in trash. Never pour unused product down the drain or on the ground.


Buyer assumes all responsibility for safety and use not in accordance with directions.

SPREADER APPLICATION PATTERN

Drop Style Spreader



Rotary Style Spreader



Net Weight: 25 Pounds

