50 ® 24-8-16

| GUARANTEED ANALYSIS: | | BY WT. |
|--------------------------------|--------|--------|
| TOTAL NITROGEN (N) | | 24.00% |
| Nitrate Nitrogen (N) | 5.00% | |
| Ammoniacal Nitrogen (N) | 5.00% | |
| Urea Nitrogen (N) | 14.00% | |
| AVAILABLE PHOSPHate (P2O5) | | 8.00% |
| SOLUBLE POTASH (K,O) | | 16.00% |
| Chlorine - not more than 1.00% | | |

STATEMENT OF SECONDARY PLANT NUTRIENTS:

| 6 |
|----|
| 6 |
| 6 |
| 6 |
| 5% |
| 16 |

SOLUBLE FERTILIZER PLUS MINORS

PRIMARY PLANT NUTRIENTS DERIVED FROM: Potassium Nitrate, Ammonium Phosphates, Urea, Ammonium Sulfate. **SECONDARY PLANT NUTRIENTS DERIVED FROM:** Ferric Sodium EDTA, Zinc Sodium EDTA, Manganous Sodium EDTA, Cupric Disodium EDTA, Sodium Borate, Sodium Molybdate.

GUARANTEED BY: Southern Agricultural Insecticides, Inc. P.O. Box 218 Palmetto, FL 34220

CAUTION: KEEP OUT OF REACH OF CHILDREN.

May be harmful if swallowed. Avoid breathing dust, and contact with skin, eyes and clothing. If ingested, give two glasses of water, induce vomiting and call a physician. If in eyes or on skin flush with water. If on clothing, launder before reuse. Wash thoroughly after handling.

Contains Potassium Nitrate, an oxidizer. Possible fire hazard.

USE PRECAUTIONS

This material is compatible with most insecticides and fungicides. Do not use in combination with Bordeaux, lime, highly alkaline materials or nutritional calcium sprays. If compatibility is not known, test on small area before use.

Timing, method and rates of application, differences in plant varieties, growing conditions and other influencing factors are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material.

Southern Agricultural Insecticides, Inc.

P.O. BOX 218
Palmetto, FL 34220
F 672
LOT NO.

P.O. BOX 429 Hendersonville, NC 28793 J2 P.O. BOX 85 Boone, NC 28607 Net Weight: 25 lbs.