CATEGOR

Range & Pasture

ACTIVE INGREDIENT:

Triclopyr 2.25 lbs AE Clopyralid .75 lbs/gal AE



PRODUCT:

Prescott Herbicide

Selective Weed Control



The premier choice for controlling broadleaf weeds including nearly all thistle and knapweed species in rangeland and permanent grass pastures as well as individual plant and spot treatments.



Range & Pasture

ACTIVE INGREDIENT:

Triclopyr 2.25 lbs AE Clopyralid .75 lbs/gal AE




PRODUCT:

Prescott Herbicide

Selective Weed Control



Alligare.com



Prescott Herbicide Use Sites

- Rangeland and permanent grass pastures
- Non-crop areas including:
 - Fence rows
 - Non-irrigation ditch banks
 - Roadsides
 - Around farm buildings
 - CRP acres

For Best Results:

Prescott Herbicide belongs to the carboxylic acid chemical family of herbicides. Herbicides in this family impact the processes of cell division, cell enlargement, protein synthesis and respiration by disrupting the hormonal balance of target plants. Prescott Herbicide is absorbed through the roots and foliage of plants, causing unregulated growth and ultimately killing target species. For best results, the spray tank solutions of Prescott Herbicide should be maintained at pH 5 to 7. Degradation of Prescott Herbicide may occur if it is used in a spray solution or with spray additives that buffer pH to below 4 or above 8.

For best activity, Prescott Herbicide must be applied when soil temperatures are above 55 degrees F and air temperatures are between 40 and 105 degrees F when plants are actively growing.

Alligare Prescott Herbicide is a broad-spectrum postemergence herbicide for control of broadleaf weeds in rangeland and permanent grass pastures, non-crop areas such as fence rows, non-irrigation ditch banks, roadsides and around farm buildings, and CRP acres. Prescott Herbicide is the premier choice for Individual Plant or Spot Treatment of many noxious and invasive weeds.

- Selective broadleaf control for the release of desirable grass species
- Non-restricted use herbicide
- Flexible tank mix options for location-specific recommendations
- Control of nearly all thistles and knapweed species

Targeted Weeds

Biennial thistles including: Ironweed, Western Poison Ivy
bull, milk, musk & plumeless Knapweed Ragweed including:
Blackberry including: diffuse, meadow, Broomweed, annual Russian & spotted Ragwort, tansy

Burdock Kudzu Skeleton, rush
Cocklebur Lettuce, prickly Sowthistle, annual and perennial
Croton Locusts Starthistle, annual

Dandelion, common and false
Dock, curly
Multiflora rose
Goldenrod
Hawkweed
Hawkweed
Horseweed
Dock, curly
Multiflora rose
Thistle, Canada
Tropical soda apple
Wormwood, absinth
Horseweed
Pigweed (suppression)

Mixing Instructions

- 1. Add 3/4 of the required spray volume to the spray tank and start agitation.
- 2. Add the required amount of Alligare Prescott Herbicide.
- 3. Add any surfactants, crop oils, or other adjuvants according to manufacturer's label.
- 4. Add any spray thickening agent, if needed to control drift, according to the manufacturer's label.
- Agitate during final filling of the spray tank and maintain sufficient agitation during application to ensure uniformity of the spray mixture.

Note: Allow time for thorough mixing of each spray ingredient before adding the next. If allowed to stand after mixing, agitate spray mixture before use.

Active Ingredient(s):

Triclopyr TEA: 3,5,6-trichloro-2-pyridinyloxyacetic acid, triethylamine salt	33.0%
Clopyralid TEA: 3,6-dichloro-2-pyridinecarboxylic acid, triethylamine salt	12.1%
OTHER INGREDIENTS:	75.6%
TOTAL:	100.0%

For more information on Prescott Herbicide or other Alligare products, or to find a regional specialist near you, call (888) 255-4427 or visit Alligare.com. Check with your Alligare regional manager for product state registration status.

©2017 Alligare LLC Always read & follow label directions.