

THURICIDE HPC

FOR THE CONTROL OF CERTAIN PLANT FEEDING CATERPILLARS AND WORMS USE 1 to 4 TEASPOONFULS PER GALLON

ACTIVE INGREDIENTS:

Sacillus thuringiensis kurstaki potency of 4,000		
International Units per milligram	0.8%	
INERTINGRE	99.2%	
TOTAL	100.0%	
Equivalent to 4.0 hillion International Linits	ner quart	

Equivalent to 4.0 billion International Units per quart. Thuricide is a registered trademark of Novartis.

KEEP OUT OF REACH OF CHILDREN CAUTION

Do not take internally. Store in a cool place in original container out of reach of children and pets.

Insect activity may be impaired by storage at temperatures above 90 degrees Fahrenheit.

Protect from freezing.

This package is intended for homeowner use.

The active ingredient concentration is the same as Thuricide packaged for use by commercial growers however commercial growers should follow label instructions on packages intended for commercial use.

CONTAINER DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS

CAUTION: Avoid breathing spray mist. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling. **FIRST AID:** If on eyes or skin, immediately flush with plenty of water. Get medical attention if irritation persists. **BUYER ASSUMES** all risks and liability from use, handling, or storage of this product not in strict accordance with the directions and precautions given herewith.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

GENERAL

Apply at first sign of infestation when worms are small and repeat at weekly intervals as needed to maintain control.

Thuricide must be carried into the insects gut while feeding, to be effective. Thorough coverage of leaf surface is essential for best results. After ingesting Thuricide, worms stop feeding and will die within 2-3 days.

Southern Agricultural Insecticides, Inc.

Palmetto, FL 34220 Hendersonville, NC 28793 Boone, NC 28607

> **EPA REG. NO 829-202** EPA EST. No. 829-FL-1

MIXING

SHAKE WELL before using - Pour recommended amount into ³/₄ filled spray tank. Agitate and maintain agitation. Do not allow dilute spray to stand in tank more than a few hours. Combinations with most fungicides and stickers are not deleterious if used promptly. A Non-ionic spreader sticker, used sparingly, is recommended when applications must be made when frequent rainfall and heavy dews are anticipated. Do not combine with alkaline materials such as bordeaux mixture or lime sulfur During hot, sunny or windy weather, late afternoon and early evening applications produce better results. Apply to obtain thorough leaf coverage.

Thuricide HPC, when applied at recommended rates, may be used up to the day of harvest.

RECOMMENDATIONS

THOROUGH COVERAGE IS ESSENTIAL. Apply indicated amount in each gallon of water and spray to obtain thorough coverage of foliage.

ORNAMENTALS

SHRUBS, TREES, AND FLOWERING PLANTS: For the control of Tent caterpillar, Fall webworm, Oakmoth larvae, Oleander Caterpillar and Gypsy moth larvae use 4 teaspoonfuls per gallon.

ORNAMENTALS IN HOME GREENHOUSES: For pests listed on this label use 2 tsp. per gallon.

CROP Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Collards, Kale, Mustard & Turnip Greens	VEGETABLES PEST Diamondback Moth, Cabbage Worm, Cabbage Looper	<u>Tsp./Ga</u> 2-4
Beans, Beets, Carrots, Celery, Chard, Endive, Escarole, Garlic, Kohlrabi, Lentils, Lettuce, Onions, Parsley, Radishes, Spinach, and Squash	Cabbage Looper	2-4
Cucumbers & Melons	Cabbage Looper	2-4
Eggplant & Peppers	Tomato Hornworm Tomato Fruitworm Cabbage Looper	1-3 2-4
Potatoes	Cabbage Looper	2-4
Tomatoes	Tomato Hornworm Cabbage Looper Tomato Fruitworm	1-3 2-4
Watermelons (FLA)	Rindworm Complex	2
	FRUITS & NUTS	-
CROP	PEST	Tsp./Ga
Almonde & Charrine	Tent Caternillar	2-4

		_
CROP Almonds & Cherries	FRUITS & NUTS PEST Tent Caterpillar	sp./Gal. 2-4
Allifolius & Chernes	Terit Caterpillar	2-4
Apples & Pears	Tufted Apple Budmoth, Variegated Leafroller, Tent Caterpillar, Fruit Tree Leafroller, and Gypsy Moth	2-4
Blueberries, Blackberries, Raspberries	Omnivorous Leafroller	2-4
Filberts	Tent Caterpillar, Filbert Leafroller	2-4
Grapes	Grape Leaf Folder	2-4
Oranges	Fruit Tree Leafroller Orange Dog Caterpillar	2-4 2
Strawberries	Cabbage Looper	2-4
Peaches & Nectarines	Tent Caterpillar, Omnivorous Leafroller	2-4
Pecans & Walnuts	Fall Webworm, Tent Caterpillar, Omnivorous leafroller	2-4