



*Home & Garden*

# CHELATED PALM NUTRITIONAL

FOR THE CORRECTION AND PREVENTION OF YELLOWING AND "FRIZZLETOP" OF PALMS  
1-2 TABLESPOONFULS PER GALLON



## DIRECTIONS FOR USE

Shake well before using.

### Palms

(For best results make both Soil and Foliar applications)

**Foliar Application:** Mix this material at the rate of 1-2 tablespoonfuls per gallon of water (1.5-3 quarts per 100 gallons). Apply to thoroughly wet the fronds to the point of runoff, also spray some of the material directly into the heart of the palm.

**Soil Application:** For small palms use 1 oz. of material in at least one half gallon of water per tree. For medium sized trees use 2-4 oz. of material in at least one gallon of water per tree. For large palms use 4-8 fluid ounces in at least 1 gallon of water per tree. Treat the root area, this area usually extends out as far as the foliage. Curative applications should be repeated in 14 days. Preventative maintenance applications should be made at 3-4 month intervals.

**NOTE:** For the continued correction and prevention of these problems the addition of SA-50 Manganese Sulfate and SA-50 Magnesium Sulfate to the soil may be advisable.

### Ornamentals, Shrubs, and Trees

**Foliage Application:** Mix this material at the rate of 1 tablespoonful per gallon of water (1.5 quarts per 100 gallons). Thoroughly cover upper and lower surfaces of foliage. Use an amount sufficient to cause runoff of spray.

**Soil Application:** Use at the rate of 1 fluid ounce (2 Tbls.) in at least 1 gallon of water per 100 sq. ft. For flower beds and vegetable gardens, treat the bed area. For shrubs and trees, treat the root area. The root area usually extends out as far as the foliage.

**NOTE:** Deficiency symptoms in many plants are an indication of improper pH. If symptoms persist or return rapidly after treatment, soil pH should be checked and adjusted if necessary.

Buyer assumes all risks of use, storage or handling not in strict accordance with the directions and precautions given herewith.

#### A micronutrient formulation for correction and prevention of secondary nutritional deficiencies

|                                       |       |
|---------------------------------------|-------|
| Soluble Manganese as (Mn).....        | 2.50% |
| Chelated Manganese as (Mn) .....      | 2.00% |
| Total Magnesium as (Mg) .....         | 1.50% |
| Water Soluble Magnesium as (Mg) ..... | 1.50% |
| Chelated Magnesium as (Mg).....       | 1.00% |
| Soluble Iron as (Fe) .....            | 1.00% |
| Chelated Iron as (Fe) .....           | 0.75% |
| Combined Sulfur as (S).....           | 3.50% |

**DERIVED FROM:** Manganese Sulfate, Manganese Lignin Sulfonate, Magnesium Sulfate, Magnesium Lignin Sulfonate, Iron Sulfate, Iron Lignin Sulfonate. Chelating agent is Lignin Sulfonate

#### USE PRECAUTIONS

THIS PRODUCT WILL SEVERELY STAIN CONCRETE, MASONRY AND PAINTED SURFACES. Be extremely careful not to allow spray or runoff to contact with these materials. Do not mix with other fertilizers, fungicides or insecticides unless compatibility is known.

## KEEP OUT OF REACH OF CHILDREN CAUTION

Avoid contact with skin and eyes.

Avoid breathing spray mist. Do not take internally.

**IF SWALLOWED:** Drink 2 glasses of water or milk and seek medical attention.

**IF IN EYES:** Flush eyes thoroughly with a large volume of water. If irritation persists seek medical attention.

**IF ON SKIN:** Rinse affected area with clear water and then wash thoroughly with soap and water.

F 672-807

# Southern Agricultural Insecticides, Inc.

Palmetto, FL 34220

Hendersonville, NC 28793

Boone, NC 28607

revised 1998