

20-20-20

SOLUBLE FERTILIZER WITH MINOR ELEMENTS

FOR USE ON ORNAMENTALS, SHRUBS, TREES,
LAWNS, HOUSE PLANTS TOBACCO,
AND VEGETABLES.

A HIGH QUALITY, WATER SOLUBLE, CONCENTRATED
FERTILIZER THAT CAN BE APPLIED AS A FOLIAR
SPRAY, SOIL DRENCH, OR WITH INJECTION RATIO
FEEDERS. THE TRACE ELEMENTS ARE CHELATED,
WHICH MAKES THEM READILY AVAILABLE TO THE
PLANT.

GUARANTEED ANALYSIS:	BY WT.
TOTAL NITROGEN (N)	20.00%
5.98% Nitrate Nitrogen	
6.00% Ammoniacal Nitrogen	
8.02% Urea Nitrogen	
AVAILABLE PHOSPHATE (P ₂ O ₅)	20.00%
SOLUBLE POTASH (K ₂ O)	20.00%
Chlorine - not more than 1.00%	

STATEMENT OF SECONDARY PLANT NUTRIENTS:

Boron as B	0.02%
Manganese as Mn (Chelated)	0.05%
Copper as Cu (Chelated)	0.05%
Molybdenum as Mo	0.0005%
Iron as Fe (Chelated)	0.10%
Zinc as Zn (Chelated)	0.05%

PRIMARY PLANT NUTRIENTS DERIVED FROM: Potassium

Nitrate, Monoammonium Phosphate, Diammonium Phosphate, Urea.

SECONDARY PLANT NUTRIENTS DERIVED FROM: Ferric Sodium
EDTA, Zinc Disodium EDTA, Manganous Disodium EDTA, Cupric
Disodium EDTA, Sodium Borate, Sodium Molybdate.

EDTA - Ethylenediaminetetraacetate ARTIFICIALLY COLORED

DRENCH APPLICATION: On plant beds and potted plants, use 2 lbs. per 100 gallons of water (2 tsp. per gallon). Apply at the rate of 1 quart per square foot or 100 gallons per 400 square feet. Repeat treatment every 1-3 weeks as needed.

CONSTANT FEED: Fertilizer applications with every watering (3 or more times per week). Use at the rate of 100-200 ppm nitrogen, 6 3/4 oz. to 13 1/2 ozs. per 100 gallons. Water with approximately 1-2 quarts per square foot or 100-200 gallons per 400 square feet (4/10 - 8/10 in. of water). Where injector systems are used, the following table may be used to determine the amount of fertilizer to use in concentrate tank. When constant feeding, it is often desirable to leach potted and benched plants about once a week in order to avoid excessive fertilizer build up, particularly on the soil surface. NOTE: Indicated rates are intended as a guide for initial applications. Experience with the particular crop and frequent soil testing are best guides to proper fertilization. (see chart)

NOTE: Do not apply with hose-end sprayers or injection ratio feeders without using a proper anti-backflow device.

LAWNS AND TURF GRASS: May be used as a supplement to regular fertilization when quick response and greening are needed. Use at the rate of 1 lb. per 10 gallons of water per 1,000 square feet. Apply monthly or as needed.

TRANSPLANT SOLUTION: For use on tobacco, vegetables such as tomato, lettuce, cabbage, cauliflower, eggplant and peppers. In general, use 3 lbs. per 100 gallons of solution (1 1/2 TBS. per gallon). Pour around plant using 1/2 to 1 pint of solution per plant. Some plants such as tobacco may be sensitive to excess amounts of fertilizer and it is best to use the low rate of 1 lb. per 100 gallons initially and check plant response. If experience dictates, up to the 3 lb. rate may be used.

This material is compatible with most insecticides and fungicides which are not highly alkaline such as Bordeaux Mixture or Lime. If compatibility is not known, test on small area before use.

Timing, method and rates of application, differences in plant varieties, growing conditions and other influencing factors are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with the directions for use and precautionary Statements on this label.

APPROXIMATE VOLUME EQUIVALENTS: 1 LB. = 2 CUPS
1/2 LB. = 1 CUP
1 oz. by weight = 1 oz. by volume (2 TBS.)

CAUTION:

KEEP OUT OF REACH OF CHILDREN.
MAY BE HARMFUL IF SWALLOWED. AVOID CONTACT
WITH EYES.

F 672

DIRECTIONS FOR USE

SPRAY APPLICATION: On Vegetables, Tobacco, Ornamentals, House Plants, Shrubs and Trees. Use 5 lbs. per 100 gallons of water (2 TBS. per gallon). Apply as a full volume spray. Repeat at 7 to 10 day intervals as needed.

OUNCES BY WEIGHT PER GALLON OF CONCENTRATE

FINAL CONC. N in ppm	INJECTOR RATIOS				
	1:50	1:100	1:128	1:150	1:200
50 ppm	1.69	3.38	4.32	5.06	6.75
75 ppm	2.53	5.06	6.48	7.59	10.13
100 ppm	3.38	6.75	8.64	10.13	13.50
125 ppm	4.22	8.44	10.80	12.66	16.88
150 ppm	5.06	10.13	12.96	15.19	20.25
175 ppm	5.90	11.81	15.12	17.72	23.63
200 ppm	6.75	13.50	17.28	20.25	27.00

Southern Agricultural Insecticides, Inc.

Palmetto, FL 34220
Hendersonville, NC 28793
Boone, NC 28607

revised 1998