

WETTABLE OR DUSTING SULFUR

ACTIVE INGREDIENT: Sulfur 90.0%
INERT INGREDIENTS: 10.0%
TOTAL: 100.0%

KEEP OUT OF REACH OF CHILDREN CAUTION

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Contact a physician.

IF INHALED: Remove victim to fresh air. Contact physician if irritation persists.

IF IN EYES: Flush eyes for 15 minutes with plenty of water, while lifting upper and lower lids. Seek medical attention if irritation persists.

IF ON SKIN: Wash all exposed skin surfaces with soap and water. Contact a physician if skin irritation develops.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: May be harmful if swallowed. Avoid inhalation of dust. Avoid contact with skin and eyes. May cause allergic response.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

1. long-sleeved shirt and long pants
2. water-proof gloves
3. shoes plus socks.

Follow manufacturers instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

For terrestrial uses, do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply when weather conditions favor drift from treated areas.

PHYSICAL OR CHEMICAL HAZARDS

When dusting, keep fire and sparks away from dust stream.

EPA EST. NO. 829-FL-1 (3); 829-NC-1 (4); 829-NC-2 (5) Number following Est. No. corresponds to first digit in Lot No.

EPA Reg. No 829-163

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box

only apply to uses of this product that are covered by the Worker Protection Standard. Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls, waterproof gloves, and shoes plus socks

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, nurseries or greenhouses. Keep children and pets of treated areas until dust has settled or sprays have dried.

BEANS as an aid in the control of rust. Apply at 4-7 day intervals using 40 lbs. per acre as a dust, or spray to obtain thorough coverage using 16 lbs. per 100 gallons (1/2 cup full per gallon). Begin applications when rust first appears and continue thereafter through harvest.

ASTER, AZALEA, CALENDULA, CHRYSANTHEMUM (hardy), CRAPE-MYRTLE, DOGWOOD, DAHLIA, DELPHINIUM, LIGUSTRUM, LILAC, MAPLE, PHLOX, RHODODENDRON, ROSE, SNAPDRAGON AND ZINNIA to control powdery mildew. Apply to obtain thorough coverage as a dust or spray thoroughly using 4 lbs. per 100 gallons (2 tablespoonfuls per gallon). Make applications at 4 to 7 day intervals beginning when powdery mildew appears and continue application as long as disease threatens.

PEACHES to control brown rot. Apply to obtain thorough coverage using 5 to 6 lbs. per 100 gallons (3 tablespoonfuls per gallon). Apply at 3 day intervals during blossom period. Apply again as soon as all petals have fallen, and repeat when 75% of the shucks have come off. Repeat applications thereafter at 10 to 14 day intervals to harvest. Use lower rate when disease presence is light. Use higher rate during periods of high rainfall and heavy disease pressure. Applications may be continued through harvest period if brown rot is present or if humid or rainy weather occurs.

CITRUS as a supplemental control of rust mite. Apply to obtain thorough coverage. Using 5 lbs. per 100 gallons (3 tablespoonfuls per gallon) as a spray or dust using 1/2 to 1 1/2 lbs. per tree depending on size of trees.

NOTE: Sulfur may cause injury to open blooms. Do not combine sulfur with oil sprays. Applications of sulfur should not be closer than 3 weeks to an oil application and this period should be extended during dry or cold weather. Applications during very hot weather 90°F or above may cause injury. Before applying this product consult proper U.S.D.A. or State Agricultural Authority for specific application procedures and dosages. Do not apply more than 50 lbs. per acre to Beans, 125 lbs. per acre to Citrus or 170 lbs. per acre to Peaches in any one application.

PROCESSING NOTE: When growing crops for processing, consult processor before applying sulfur.

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Store pesticide in the original container in a locked storage area.

PESTICIDE DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash. **CONTAINER DISPOSAL:** Do not reuse bag. Discard bag in trash. **NOTICE:** Buyer assumes all risks of use, storage, or handling of this material not in strict accordance with DIRECTIONS FOR USE and PRECAUTIONARY STATEMENTS stated on this label.

Southern Agricultural Insecticides, Inc.

Palmetto, FL 34220 Hendersonville, NC 28793

Boone, NC 28607

revised 1998